Product Name: Recombinant Mouse FLT3LG (C-6His)

Catalog #: PHM0677



Summary

Name Flt3L/Flt3-Ligand/Fms-related tyrosine kinase 3 ligand

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <0.05EU/μg as determined by LAL test.

Construction Recombinant Mouse Fms-like Tyrosine Kinase 3 Ligand is produced by our

Mammalian expression system and the target gene encoding Gly27-Arg188 is

expressed with a 6His tag at the C-terminus.

Accession # P49772

Host Human Cells

Species Mouse

Predicted Molecular Mass 19.4 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Stability&Storage Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt.

Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at ≤ -20°C for 3 months.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is


not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Product Name: Recombinant Mouse FLT3LG (C-6His)

Catalog #: PHM0677





Alternative Names

Fms-related tyrosine kinase 3 ligand (Flt3L); SL cytokine; Flt3 ligand;

Background

Fms-related tyrosine kinase 3 ligand(Flt3L) is a single-pass type I membrane protein and consists of 232 amino acids. Flt3L is a hematopoietic four helical bundle cytokine, structurally homologous to stem cell factor and colony stimulating factor. Flt3L synergizes well with a number of other colony stimulating factors and interleukins. Flt3L stimulates the proliferation and differentiation of various blood cell progentiors by activating FLT3.

Note

For Research Use Only, Not for Diagnostic Use.