


Summary

Name	Flt3L/Flt3-Ligand/Fms-related tyrosine kinase 3 ligand
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Mouse Fms-like Tyrosine Kinase 3 Ligand is produced by our Mammalian expression system and the target gene encoding Gly27-Arg188 is expressed with a human IgG1 Fc tag at the C-terminus.
Accession #	P49772
Host	Human Cells
Species	Mouse
Predicted Molecular Mass	45.5 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Product Name: Recombinant Mouse FLT3LG (C-Fc)
Catalog #: PHM0678



Alternative Names

Fms-related tyrosine kinase 3 ligand (Flt3L); SL cytokine; Flt3 ligand

Background

Fms-related tyrosine kinase 3 ligand(Flt3L) is a single-pass type I membrane protein and consists of 232 amino acids. Flt3L is a hematopoietic four helical bundle cytokine, structurally homologous to stem cell factor and colony stimulating factor. Flt3L synergizes well with a number of other colony stimulating factors and interleukins. Flt3L stimulates the proliferation and differentiation of various blood cell progenitors by activating FLT3.

Note

For Research Use Only , Not for Diagnostic Use.