

**Product Name: Recombinant Mouse CD74 (C-6His)**  
**Catalog #: PHM0795**




---

## Summary

<b>Name</b>	CD74/HLADG/H-2 class II histocompatibility antigen gamma chain
<b>Purity</b>	Greater than 95% as determined by reducing SDS-PAGE
<b>Endotoxin level</b>	<1 EU/μg as determined by LAL test.
<b>Construction</b>	Recombinant Mouse HLA Class II Histocompatibility Antigen Gamma Chain is produced by our Mammalian expression system and the target gene encoding Gln56-Leu215 is expressed with a 6His tag at the C-terminus.
<b>Accession #</b>	P04441-2
<b>Host</b>	Human Cells
<b>Species</b>	Mouse
<b>Predicted Molecular Mass</b>	19.4 KDa
<b>Formulation</b>	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
<b>Stability&amp;Storage</b>	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
<b>Reconstitution</b>	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## SDS-PAGE image

**Product Name: Recombinant Mouse CD74 (C-6His)**  
**Catalog #: PHM0795**



### Alternative Names

Cluster of Differentiation 74;CD74 antigen; CD74 molecule, major histocompatibility complex, class II invariant chain; DHLAG gamma chain of class II antigens; HLA class II histocompatibility antigen gamma chain; HLADG; HLA-DR antigens-associated invariant chain; HLA-DR-gamma; Ia antigen-associated invariant chain; Ia-associated invariant chain; Ia-GAMMA; MHC HLA-DR gamma chain; CD74; DHLAG; HLADG; Ia-gamma; INVG34;

### Background

Mouse HLA class II histocompatibility antigen gamma chain (CD74), is a single-pass type II membrane glycoprotein which contains 1 thyroglobulin type-1 domain. Mouse CD74 shares 75% and 88% aa sequence identity with human and rat CD74, respectively. CD74 plays an important role in adaptive immunity, inflammation, and cancer. It plays a critical role in MHC class II antigen processing by stabilizing peptide-free class II alpha/beta heterodimers in a complex soon after their synthesis and directing transport of the complex from the endoplasmic reticulum to compartments where peptide loading of class II takes place. CD74 also associates with CD44 and binds with high affinity to the cytokine MIF, leading to inflammatory leukocyte responses, protection from tissue fibrosis, B cell proliferative and survival signaling, and the up-regulation of angiogenic factors in endometrial stromal cells.

### Note

For Research Use Only , Not for Diagnostic Use.