


Summary

Name	HVEM/TNFRSF14/CD270
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/µg as determined by LAL test.
Construction	Recombinant Mouse Herpesvirus Entry Mediator is produced by our Mammalian expression system and the target gene encoding Gln39-Val207 is expressed with a human IgG1 Fc tag at the C-terminus.
Accession #	Q80WM9
Host	Human Cells
Species	Mouse
Predicted Molecular Mass	45.47 KDa
Formulation	Lyophilized from a 0.2 μ m filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background



Alternative NamesTnfrsf14; Herpesvirus entry mediator; HVEM; TR2; TNF receptor-like molecule;
ATAR; another TRAF-associated receptor; Tumor necrosis factor receptor
superfamily member 14BackgroundMouse Protein Tnfrsf14, is a type I transmembrane protein belonging to the TNF
receptor superfamily. It is tumor necrosis factor receptor superfamily member 14
and expressed on the surface of T cells during the resting state. Interaction of
HVEM with TNF family member LIGHT co-stimulates T cells and promotes
inflammation. HVEM also triggers inhibitory signaling cascade in effector T (Teff)
cells and regulatory T cells (Tregs) as a ligand of B and T lymphocyte attenuator.
Tnfrsf14 is detected in peripheral blood T cells, B cells, monocytes and in various
tissues enriched in lymphoid cells. It has demonstrated that HVEM Ig is able to
exert a significant antiviral effect against HSV-1 infection in vivo.

Note

For Research Use Only, Not for Diagnostic Use.