

Product Name: Recombinant Mouse IGFBP-7 (C-6His)
Catalog #: PHM0846




Summary

Name	IGFBP-7/Igfbp7
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Mouse Insulin-like Growth Factor-binding Protein 7 is produced by our Mammalian expression system and the target gene encoding Ser26-Leu281 is expressed with a 6His tag at the C-terminus.
Accession #	Q61581
Host	Human Cells
Species	Mouse
Predicted Molecular Mass	27.2 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, 150mM NaCl, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Product Name: Recombinant Mouse IGFBP-7 (C-6His)
Catalog #: PHM0846



Alternative Names

Insulin-like growth factor-binding protein 7;IGFBP7;IGF-binding protein 7;IGFBP-rP1;MAC25 protein;Tumor-derived adhesion factor;TAF

Background

Insulin-like growth factor-binding protein 7(IGFBP-7) is a secreted glycosylated protein that contains three protein domain modules. IGFBP7 contains an N-terminal IGFBP domain, followed by a Kazal-type serine proteinase inhibitor domain and a C-terminal immunoglobulin-like C2-type domain. Human and mouse IGFBP7 are highly homologous and share 94% aa sequence identity. It is expressed in many normal tissues and in cancer cells. It is abundantly expressed in high endothelial venules (HEVs) of blood vessels in the secondary lymphoid tissues. It binds IGF and insulin with very low affinity and has been shown to enhance the mitogenic actions of IGF and insulin. IGFBP7 also has IGF/insulin-independent activities. It interacts with heparan sulfate proteoglycans, type IV collagen, and specific chemokines. It supports weak cell adhesion, promotes cell spreading on type IV collagen, and stimulates the production of the potent vasodilator PGI₂. It modulates tumor cell growth and has also been implicated in angiogenesis.

Note

For Research Use Only , Not for Diagnostic Use.