


Product Name: Recombinant Mouse IL-2RA (C-6His)
Catalog #: PHM0914



Summary

Name	IL-2 R alpha/CD25/IL2-RA/IL-2R α /IL-2 receptor subunit alpha
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/ μ g as determined by LAL test.
Construction	Recombinant Mouse Interleukin-2 Receptor Subunit Alpha is produced by our Mammalian expression system and the target gene encoding Glu22-Lys236 is expressed with a 6His tag at the C-terminus.
Accession #	P01590
Host	Human Cells
Species	Mouse
Predicted Molecular Mass	25.5 KDa
Formulation	Lyophilized from a 0.2 μ m filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at $\leq -70^{\circ}\text{C}$, stable for 6 months after receipt. Store at $\leq -70^{\circ}\text{C}$, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μ g/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Mouse IL-2RA (C-6His)
Catalog #: PHM0914



Alternative Names

CD25; IL-2 R alpha; IL2RA; CD25 antigen; IDDM10; IL-2 receptor subunit alpha; IL2R; IL-2R subunit alpha; IL-2-RA; IL2-RA; interleukin 2 receptor; alpha; interleukin-2 receptor subunit alpha; p55; TAC antigen; TCGFR

Background

The interleukin 2 receptor alpha (IL2RA), also called CD25, is a type I transmembrane protein that presents on activated T cells, activated B cells, some thymocytes, myeloid precursors, and oligodendrocytes. IL2RA is a member of cytokine receptors family that utilizes the common gamma chain subunit (γc). IL2RA is expressed in most B-cell neoplasms, some acute nonlymphocytic leukemias, neuroblastomas, and tumor infiltrating lymphocytes. IL2RA associates with IL2RB (CD122) to form a heterodimer that can act as a high-affinity receptor for IL-2.

Note

For Research Use Only , Not for Diagnostic Use.