

**Product Name: Recombinant Mouse IL-1R1 (C-Fc)**  
**Catalog #: PHM0975**




---

## Summary

<b>Name</b>	IL-1RI/Interleukin-1 receptor type 1/IL-1R1
<b>Purity</b>	Greater than 95% as determined by reducing SDS-PAGE
<b>Endotoxin level</b>	<1 EU/μg as determined by LAL test.
<b>Construction</b>	Recombinant Mouse Interleukin-1 Receptor Type 1 is produced by our Mammalian expression system and the target gene encoding Leu20-Lys338 is expressed with a human IgG1 Fc tag at the C-terminus.
<b>Accession #</b>	P13504
<b>Host</b>	Human Cells
<b>Species</b>	Mouse
<b>Predicted Molecular Mass</b>	64 KDa
<b>Formulation</b>	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
<b>Stability&amp;Storage</b>	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
<b>Reconstitution</b>	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## SDS-PAGE image

**Product Name: Recombinant Mouse IL-1R1 (C-Fc)**  
**Catalog #: PHM0975**



### **Alternative Names**

Interleukin-1 receptor type 1; IL-1R-1; IL-1RT-1; IL-1RT1; CD121 antigen-like family member A; Interleukin-1 receptor alpha; IL-1R-alpha; p80; CD121a; mL-1R1

### **Background**

Mouse Interleukin-1 receptor type 1/IL-1 RI is a cytokine receptor that belongs to the interleukin-1 receptor family. This protein is a receptor for interleukin 1 alpha (IL1A), interleukin 1 beta (IL1B), and interleukin 1 receptor antagonist (IL1RA). It is an important mediator involved in many cytokine induced immune and inflammatory responses. An IL1 receptor accessory protein that can heterodimerize with the Type I receptor in the presence of IL1 $\alpha$  or IL1 $\beta$  but not IL1ra, was identified. This Type I receptor complex appears to mediate all the known IL1 biological responses. The receptor Type II has a short cytoplasmic domain and does not transduce IL1 signals. In addition to the membranebound form of IL1 RII, a naturally occurring soluble form of IL1 RII has been described. It has been suggested that the Type II receptor, either as the membranebound or as the soluble form, serves as a decoy for IL1 and inhibits IL1 action by blocking the binding of IL1 to the signaling Type I receptor complex.

### **Note**

For Research Use Only , Not for Diagnostic Use.