

Product Name: Recombinant Mouse ST2 (V192A, C-Fc)
Catalog #: PHM0982




Summary

Name	sST2/IL-33 R/Interleukin-1 Receptor-Like 1/IL-1RL1/IL-1 R4/DER4 (V192A)
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Mouse Interleukin-1 receptor-like 1 is produced by our Mammalian expression system and the target gene encoding Ser27-Ala337 (Val192Ala) is expressed with a human IgG1 Fc tag at the C-terminus.
Accession #	P14719-2
Host	Human Cells
Species	Mouse
Predicted Molecular Mass	62.7 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Product Name: Recombinant Mouse ST2 (V192A, C-Fc)
Catalog #: PHM0982



Alternative Names

Interleukin-1 receptor-like 1; Lymphocyte antigen 84; Protein ST2; Protein T1; Il1rl1; DER4; Fit-1; IL-1 R4; IL-1RL1; IL-33R; Ly84; ST2L; ST2V; suppression of tumorigenicity 2

Background

ST2, also called IL-1 R4, is an Interleukin-1 receptor family glycoprotein that plays a role in Th2 immune responses. ST2 is expressed on the surface of mast cells, activated Th2 cells, macrophages, and cardiac myocytes. This receptor is very similar to the IL-1 receptor type I and the IL-18 receptor α chain in that ST2 also has three extracellular Ig domains and an intracellular Toll domain. ST2 binds IL-33, enhances inflammatory cytokines by activating nuclear factor- κ B (NF- κ B) and mitogen activated protein (MAP) kinases. ST2 exists as either a membrane bound form (ST2L) or as a soluble form (sST2). ST2L acts as a transmembrane signalling receptor for IL-33 by mediating the effect of IL-33 on the inflammatory process, while sST2 can suppress IL-33 activity.

Note

For Research Use Only , Not for Diagnostic Use.