


Summary

Name	LAIR1/CD305
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Mouse Leukocyte-associated Immunoglobulin-like Receptor 1 is produced by our Mammalian expression system and the target gene encoding Gln22-Tyr141 is expressed with a 6His tag at the C-terminus.
Accession #	Q8BG84
Host	Human Cells
Species	Mouse
Predicted Molecular Mass	14.4 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Product Name: Recombinant Mouse LAIR1 (C-6His)
Catalog #: PHM1060



Alternative Names

Leukocyte-associated immunoglobulin-like receptor 1; LAIR-1; mLAIR-1; CD305; Lair1

Background

Leukocyte-associated Ig-like receptor-1 (LAIR-1) is an inhibitory receptor of the Ig superfamily that is structurally related to inhibitory members of KIR and ILT/CD85 families. It is expressed on immune cells, including NK cells, T cells, B cells, monocytes, immature neutrophils, dendritic cells and most thymocytes. The 253 amino acid (aa) type I transmembrane (TM) protein contains a 21 aa signal sequence, a 124 aa extracellular domain (ECD), a 20 aa TM domain and a 98 aa cytoplasmic domain. The ECD includes one C2-type Ig-like domain and two potential N-linked glycosylation sites. Tyrosine phosphorylation of two cytoplasmic ITIM motifs results in recruitment of phosphatases and down-regulation of signaling through activating receptors. LAIR1 shows high-affinity binding of collagens that results in inhibition of degranulation in a basophilic leukemia cell line.

Note

For Research Use Only , Not for Diagnostic Use.