


## Summary

|                                 |  |
|---------------------------------|--|
| <b>Name</b>                     | IGFBP-9/Nephroblastoma Overexpressed Gene/NOV/CCN3   |
| <b>Purity</b>                   | Greater than 95% as determined by reducing SDS-PAGE  |
| <b>Endotoxin level</b>          | <1 EU/μg as determined by LAL test.  |
| <b>Construction</b>             | Recombinant Mouse Nephroblastoma Overexpressed Gene is produced by our Mammalian expression system and the target gene encoding Ser26-Ile354 is expressed with a 6His tag at the C-terminus.   |
| <b>Accession #</b>              | Q64299   |
| <b>Host</b>                     | Human Cells  |
| <b>Species</b>                  | Mouse  |
| <b>Predicted Molecular Mass</b> | 37.1 KDa   |
| <b>Formulation</b>              | Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.  |
| <b>Shipping</b>                 | The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.   |
| <b>Stability&amp;Storage</b>    | Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.   |
| <b>Reconstitution</b>           | Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. |

## SDS-PAGE image

**Product Name: Recombinant Mouse NOV (C-6His)**  
**Catalog #: PHM1204**



### Alternative Names

Protein NOV homolog; NovH; CCN family member 3; Nephroblastoma-overexpressed gene protein homolog; Nov

### Background

NOV, also called CCN3, is a secreted protein of CCN family members. CCN family members are highly conserved cysteine rich proteins sharing a common modular structure having 4 conserved domains, insulin-like growth factor-binding protein (IGFBP) domain, von Willebrand type C (VWC) domain, thrombospondin-1 (TSP-1) domain, and C-terminal (CT) domain (absent in CCN5). By specific interactions with these domains, CCN proteins modulate multiple signalling pathways including BMPs, Wnt, TGFs, Notch and integrins to regulate cell proliferation, differentiation, adhesion, migration, angiogenesis, and survival. CCN3 is firstly characterized as a promoter of progenitor activity of human hematopoietic stem cells, as knockdown of CCN3 can abrogate the function of primitive progenitors. Recent studies showed that CCN3 is also actively involved in the process of wound healing. CCN3 is highly expressed in granulation tissues of cutaneous wounds and capable of inducing synthetic responses of fibroblasts.

### Note

For Research Use Only , Not for Diagnostic Use.