


Product Name: Recombinant Mouse PDGFRA (C-6His)
Catalog #: PHM1288



Summary

Name	PDGF R alpha/PDGFRA/CD140a
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Mouse Platelet-derived Growth Factor Receptor Alpha is produced by our Mammalian expression system and the target gene encoding Leu25-Glu524 is expressed with a 6His tag at the C-terminus.
Accession #	P26618
Host	Human Cells
Species	Mouse
Predicted Molecular Mass	56.9 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Mouse PDGFRA (C-6His)
Catalog #: PHM1288



Alternative Names

Platelet-derived growth factor receptor alpha; PDGF-R-alpha; Alpha platelet-derived growth factor receptor; Alpha-type platelet-derived growth factor receptor; CD140a; Pdgfra

Background

Platelet-derived growth factor receptors (PDGFR) are cell surface tyrosine kinase receptors for members of the platelet-derived growth factor (PDGF) family. The PDGF family consists of PDGF-A, -B, -C and -D, which form either homo- or heterodimers (PDGF-AA, -AB, -BB, -CC, -DD). The four PDGFs are inactive in their monomeric forms. PDGFs bind to the protein tyrosine kinase receptors PDGF receptor- α and - β . These two receptor isoforms dimerize upon binding the PDGF dimer, leading to three possible receptor combinations, namely - $\alpha\alpha$, - $\beta\beta$ and - $\alpha\beta$. PDGFR α and PDGFR β are members of the class III RTK family. Inappropriate PDGFR α and PDGFR β signaling has been linked to a number of proliferative disorders.

Note

For Research Use Only , Not for Diagnostic Use.