Product Name: Recombinant Mouse PLGF (C-6His)

Catalog #: PHM1319



Summary

Name Placenta growth factor/PIGF/PGF/PLGF

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/μg as determined by LAL test.

Construction Recombinant Mouse Placenta Growth Factor is produced by our Mammalian

expression system and the target gene encoding Val19-Pro158 is expressed

with a 6His tag at the C-terminus.

Accession # P49764

Host Human Cells

Species Mouse

Predicted Molecular Mass 16.9 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Stability&Storage Lyophilized protein should be stored at \leq -20°C, stable for one year after receipt.


Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at \leq -20°C for 3 months.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Mouse PLGF (C-6His) Catalog #: PHM1319

C EnkiLife

Alternative Names Placenta growth factor; PIGF; PIgf; D12S1900; PGFL; PGF; PIGF-2; SHGC-10760

Background Placental growth factor is a protein that in humans is encoded by the PGF gene. It

is a secreted protein and belongs to the PDGF/VEGF growth factor family. The protein is a member of the VEGF (vascular endothelial growth factor) sub-family-a key molecule in angiogenesis and vasculogenesis, in particular during embryogenesis. The main source of PGF during pregnancy is the placental trophoblast. PGF is also expressed in many other tissues, including the villous

trophoblast.

Note

For Research Use Only, Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838