


Product Name: Recombinant Mouse CD31 (C-6His)
Catalog #: PHM1336



Summary

Name	CD31/PECAM-1/Platelet Endothelial Cell Adhesion Molecule
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Mouse Platelet Endothelial Cell Adhesion Molecule is produced by our Mammalian expression system and the target gene encoding Glu18-Lys590 is expressed with a 6His tag at the C-terminus.
Accession #	Q08481
Host	Human Cells
Species	Mouse
Predicted Molecular Mass	63.4 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, 5mM EDTA, pH7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Product Name: Recombinant Mouse CD31 (C-6His)
Catalog #: PHM1336



Background

Alternative Names Platelet endothelial cell adhesion molecule; PECAM-1; CD31; Pecam1; Pecam

Background Platelet endothelial cell adhesion molecule (PECAM-1, CD31) is a type I transmembrane glycoprotein adhesion molecule in the immunoglobulin superfamily. PECAM-1 is concentrated at cell junctions and is required for transendothelial migration (TEM). The extracellular domain (ECD) of PECAM-1 has ten potential N-linked glycosylation sites and six C2-type Ig-like domains, the first of which is critical for adhesion and extravasation. The cytoplasmic domain contains immunoregulatory tyrosine-based inhibitory and switch motifs (ITIM, ITSM) that mediate both inhibition and activation via phosphotyrosine-mediated engagement of SH2-containing signaling molecules. Expression is restricted to cells involved in circulation, especially endothelial cells, platelets, monocytes, neutrophils and lymphocyte subsets. PECAM-1 participates with other adhesion molecules in some functions, but is the critical molecule for TEM. Homotypic PECAM-1 adhesion in trans, combined with cycling of PECAM-1 to and from surface-connected endothelial cell vesicles, leads leukocytes across endothelial tight junctions.

Note

For Research Use Only , Not for Diagnostic Use.