


Summary

Name	VISTA/B7-H5/Gi24/C10orf54/PD-1H/platelet receptor Gi24
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Mouse Platelet Receptor Gi24 is produced by our Mammalian expression system and the target gene encoding Phe33-Ala191 is expressed with a 6His tag at the C-terminus.
Accession #	Q9D659
Host	Human Cells
Species	Mouse
Predicted Molecular Mass	18.6 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Mouse VISTA (C-6His)
Catalog #: PHM1339



Alternative Names

Platelet receptor Gi24; stress induced secreted protein 1; Dies1; VISTA; SISP1; B7-H5; PD-1H;Gi24

Background

Mouse Platelet receptor Gi24(VISTA) is a transmembrane glycoprotein with homology to B7like immune costimulatory molecules. Mature mouse Gi24 contains a 159 amino acid (aa) extracellular domain (ECD) with one V-type Ig-like domain, a 21 aa transmembrane segment, and a 97 aa cytoplasmic domain. VISTA promotes both MT1-MMP expression and the MT1-MMP mediated activation of MMP-2. It supports the differentiation of embryonic stem cells (ESC) and enhances BMP-4 induced signaling in ESC, but it is also down-regulated following BMP-4 exposure. It binds to BMP-4 directly and also associates with the type I BMP receptor Activin RIB/ALK-4. It is expressed on the surface of naïve CD4+ T cells and regulatory T cells. It is up-regulated in vivo on activated monocytes and dendritic cells. VISTA inhibits CD4+ and CD8+ T cell proliferation and their production of IL-2 and IFN- γ . Its expression on tumor cells attenuates the antitumor immune response and enables more rapid tumor progression.

Note

For Research Use Only , Not for Diagnostic Use.