

**Product Name: Recombinant Mouse CREG (C-6His)**  
**Catalog #: PHM1382**




## Summary

<b>Name</b>	CREG/CREG1/Protein CREG1
<b>Purity</b>	Greater than 95% as determined by reducing SDS-PAGE
<b>Endotoxin level</b>	<1 EU/μg as determined by LAL test.
<b>Construction</b>	Recombinant Mouse Cellular Repressor Of E1A-stimulated Genes is produced by our Mammalian expression system and the target gene encoding Arg32-Gln220 is expressed with a 6His tag at the C-terminus.
<b>Accession #</b>	O88668
<b>Host</b>	Human Cells
<b>Species</b>	Mouse
<b>Predicted Molecular Mass</b>	22.5 KDa
<b>Formulation</b>	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
<b>Stability&amp;Storage</b>	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
<b>Reconstitution</b>	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## SDS-PAGE image

**Product Name: Recombinant Mouse CREG (C-6His)**  
**Catalog #: PHM1382**



### **Alternative Names**

CREG1; cellular repressor of E1A-stimulated genes; cellular repressor of E1A-stimulated genes 1CREG; protein CREG1

### **Background**

Cellular repressor of E1A genes (CREG) is an evolutionarily conserved lysosomal protein, and an important new factor in regulating tissue homeostasis that has been shown to antagonize injury of tissues or cells. CREG contains three mannose 6-phosphate (M6P) markers, and depends on interactions with M6P receptors for efficient delivery to lysosomes, which is implicated in the regulation of lysosomal functions. This protein shares limited sequence similarity with E1A and binds both the general transcription factor TBP and the tumor suppressor pRb in vitro. CREG plays an important role in the control of cell growth and differentiation. It has been shown that CREG antagonizes transcriptional and cellular transformation by the adenoviral E1A oncoprotein, induces differentiation while attenuating cellular proliferation, regulates the levels of the signaling kinases ERK1/2, and mediates glucocorticoid-induced proliferation of ileal epithelial cells. CREG is widely expressed in adult tissues, such as the brain, heart, lungs, liver, intestines and kidneys in mice, but is not markedly expressed in pluripotent embryonic stem cells.

### **Note**

For Research Use Only , Not for Diagnostic Use.