


Product Name: Recombinant Mouse TIGIT (C-Fc)
Catalog #: PHM1608



Summary

| | |
|---------------------------------|--|
| Name | TIGIT/VSIG9/VSTM3/T-cell immunoreceptor with Ig and ITIM domains |
| Purity | Greater than 95% as determined by reducing SDS-PAGE |
| Endotoxin level | <1 EU/μg as determined by LAL test. |
| Construction | Recombinant Mouse T-cell Immunoreceptor With Ig Wnd ITIM Domains is produced by our Mammalian expression system and the target gene encoding Gly26 - Thr143 is expressed with a human IgG1 Fc tag at the C-terminus. |
| Accession # | NP_001139797 |
| Host | Human Cells |
| Species | Mouse |
| Predicted Molecular Mass | 40.1 KDa |
| Formulation | Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4. |
| Shipping | The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below. |
| Stability&Storage | Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles. |
| Reconstitution | Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. |

SDS-PAGE image



Background

Product Name: Recombinant Mouse TIGIT (C-Fc)
Catalog #: PHM1608



Alternative Names

T-cell immunoreceptor with Ig and ITIM domains; Tigit

Background

T cell immunoreceptor with Ig and ITIM domains (TIGIT), also called WUCAM, VSIG9 and Vstm3, is a member of the CD28 family within the Ig superfamily of proteins. TIGIT contains an immunoglobulin variable domain, a transmembrane domain and an immunoreceptor tyrosine-based inhibitory motif (ITIM), and is expressed on regulatory, memory, activated T cells and NK cells. TIGIT binds to CD155(PVR) that appear on dendritic cells (DC), macrophages and endothelium with high affinity, and CD112(PVRL2) with lower affinity, but not CD113 (PVRL3). TIGIT-Fc fusion protein could interact with PVR on DC and enhance the secretion of IL-10, but inhibit the macrophage activation. Mice lacking TIGIT show increased T cell responses and susceptibility to autoimmune challenges, while knockdown of TIGIT with siRNA in human memory T cells did not affect T cell responses.

Note

For Research Use Only , Not for Diagnostic Use.