


Product Name: Recombinant Mouse CD96 (C-6His)
Catalog #: PHM1611



Summary

Name	CD96/T-cell Surface Protein Tactile
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Mouse T-cell Surface Protein Tactile is produced by our Mammalian expression system and the target gene encoding Val22-Met536 is expressed with a 6His tag at the C-terminus.
Accession #	Q3U0X8
Host	Human Cells
Species	Mouse
Predicted Molecular Mass	57.6 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Product Name: Recombinant Mouse CD96 (C-6His)
Catalog #: PHM1611



Background

Alternative Names

T-cell surface protein tactile; Cell surface antigen CD96; T cell-activated increased late expression protein; CD96

Background

The cluster of differentiation (CD) system is commonly used as cell markers in immunophenotyping. Different kinds of cells in the immune system can be identified through the surface CD molecules which associating with the immune function of the cell. The CD155 ligand CD96 is a member of the Ig superfamily. Its a immunoglobulin-like protein tentatively allocated to the repertoire of human NK receptors. NK cells recognize poliovirus receptor (PVR), anectins and nectin-like protein family member serve to mediate cell-cell adhesion, cell migration, with the presence of an additional receptor, CD96. CD96 promotes NK cell adhesion to target cells expressing PVR, stimulates cytotoxicity of activated NK cells, and mediates acquisition of PVR from target cells.

Note

For Research Use Only , Not for Diagnostic Use.