


Summary

Name	CD28/TP44/T-cell-specific surface glycoprotein CD28
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/µg as determined by LAL test.
Construction	Recombinant Mouse T-cell-specific Surface Glycoprotein CD28 is produced by our Mammalian expression system and the target gene encoding Asn20- Lys149 is expressed with a human IgG1 Fc, 6His tag at the C-terminus.
Accession #	P31041
Host	Human Cells
Species	Mouse
Predicted Molecular Mass	43 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background



Alternative Names T-cell-specific surface glycoprotein CD28;CD28

Background T-cell-specific surface glycoprotein CD28, contains an Ig-like V-type domain. CD28 is one of the proteins expressed on T cells that provide co-stimulatory signals, which are required for their activation. It is the receptor for CD80 and CD86. When activated by Toll-like receptor ligands, the CD80 expression is upregulated in antigen presenting cells (APCs). The CD86 expression on antigen presenting cells is constitutive. CD28 is the only B7 receptor constitutively expressed on naive T cells. It involved in T-cell activation, the induction of cell proliferation and cytokine production and promotion of T-cell survival.

Note

For Research Use Only, Not for Diagnostic Use.