


**Product Name: Recombinant Mouse TSLP (C-Fc)**  
**Catalog #: PHM1640**



## Summary

|                                 |  |
|---------------------------------|--|
| <b>Name</b>                     | TSLP/Thymic stromal lymphopoietin  |
| <b>Purity</b>                   | Greater than 95% as determined by reducing SDS-PAGE  |
| <b>Endotoxin level</b>          | <1 EU/μg as determined by LAL test.  |
| <b>Construction</b>             | Recombinant Mouse Thymic Stromal Lymphopoietin is produced by our Mammalian expression system and the target gene encoding Tyr20-Glu140 is expressed with a human IgG1 Fc tag at the C-terminus.   |
| <b>Accession #</b>              | Q9JIE6   |
| <b>Host</b>                     | Human Cells  |
| <b>Species</b>                  | Mouse  |
| <b>Predicted Molecular Mass</b> | 41.1 KDa   |
| <b>Formulation</b>              | Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.  |
| <b>Shipping</b>                 | The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.   |
| <b>Stability&amp;Storage</b>    | Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.   |
| <b>Reconstitution</b>           | Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. |

## SDS-PAGE image



**Product Name: Recombinant Mouse TSLP (C-Fc)**  
**Catalog #: PHM1640**

---



## Background

**Alternative Names** Thymic stromal lymphopoietin;Thymic stroma-derived lymphopoietin;Tslp

**Background** Thymic stromal lymphopoietin (TSLP) is a protein belonging to the cytokine family, contains 140 amino acids. It is known to play an important role in the maturation of T cell populations through activation of antigen presenting cells. TSLP induces the release of T-cell-attracting chemokines from monocytes and, in particular, enhances the maturation of CD11c+ dendritic cells. It can induce allergic inflammation by directly activating mast cells. TSLP is produced mainly by non-hematopoietic cells such as fibroblasts, epithelial cells and different types of stromal or stromal-like cells. These cells are located in regions where TSLP activity is required.

## Note

For Research Use Only , Not for Diagnostic Use.