

Product Name: Recombinant Mouse TREM-1 (C-6His)
Catalog #: PHM1717




Summary

Name	TREM-1/CD354/Triggering receptor expressed on myeloid cells 1
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Mouse Triggering Receptor Expressed On Myeloid Cells 1 is produced by our Mammalian expression system and the target gene encoding Ala21-Ser202 is expressed with a 6His tag at the C-terminus.
Accession #	Q9JKE2
Host	Human Cells
Species	Mouse
Predicted Molecular Mass	20.9 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Product Name: Recombinant Mouse TREM-1 (C-6His)
Catalog #: PHM1717



Alternative Names

Triggering receptor expressed on myeloid cells 1; TREM-1; CD354; Trem1

Background

Triggering Receptor Expressed on Myeloid Cells 1 (TREM-1) is a transmembrane protein with a single Ig-like domain. TREM-1 associates with the adapter protein, DAP12, to deliver an activating signal. TREM-1 is expressed on blood neutrophils and monocytes, and the expression is up-regulated by bacterial LPS. TREM-1 is expressed at high levels on neutrophils of patients with microbial sepsis and in mice with a TREM-1/Fc fusion protein protected mice against LPS-induced shock. It stimulates neutrophil and monocyte-mediated inflammatory responses. Triggers release of pro-inflammatory chemokines and cytokines, as well as increased surface expression of cell activation markers. TREM-1 is amplifier of inflammatory responses that are triggered by bacterial and fungal infections and are a crucial mediator of septic shock.

Note

For Research Use Only , Not for Diagnostic Use.