

Product Name: Recombinant Mouse TREM-2b (C-Fc)
Catalog #: PHM1724




Summary

Name	TREM-2b/Triggering Receptor Expressed on Myeloid Cells 2b
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Mouse Triggering Receptor Expressed On Myeloid Cells 2b is produced by our Mammalian expression system and the target gene encoding Leu19-Pro168 is expressed with a human IgG1 Fc tag at the C-terminus.
Accession #	Q99NH8
Host	Human Cells
Species	Mouse
Predicted Molecular Mass	43.7 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Product Name: Recombinant Mouse TREM-2b (C-Fc)
Catalog #: PHM1724



Alternative Names

Triggering Receptor Expressed on Myeloid Cells 2b; Triggering receptor expressed on myeloid cells 2; TREM-2; Triggering receptor expressed on monocytes 2; Trem2; Trem2a; Trem2b; Trem2c; TREM-2b

Background

Triggering receptor expressed on myeloid cells-2 (TREM-2) is a cell surface receptor primarily expressed on macrophages, osteoclasts, microglia and dendritic cells. TREM-2 is one member of the TREM family, inhibiting the releasing of inflammatory mediators, so it is an important in vivo anti-inflammatory receptor. TREM-2 consists of an 18 aa signal sequence, a 153 aa extracellular domain (ECD) with one V-type Ig-like domain, a 21 aa transmembrane (TM) domain, and a 35 aa cytoplasmic tail. A soluble form of TREM-2 (TREM-2b) created by alternate splicing diverges at aa 161. TREM-2 transduces intracellular signals through the adaptor DAP12. After binding of TREM-2 with ligand, the TREM-2/DAP12 (dead-cell-activated-receptor-associated protein)-mediated signal transduction pathway causes a series of intracellular protein tyrosine phosphorylation reactions and enzymatic reactions, which then activate the myeloid cells and participate T cell responses.

Note

For Research Use Only , Not for Diagnostic Use.