


Product Name: Recombinant Mouse OX40L (N-8His)
Catalog #: PHM1742



Summary

Name	OX40 Ligand/OX40L/TNFSF4/CD252/Tumor necrosis factor ligand superfamily member 4
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Mouse OX40 Ligand is produced by our Mammalian expression system and the target gene encoding Ser51-Leu198 is expressed with a 8His tag at the N-terminus.
Accession #	P43488
Host	Human Cells
Species	Mouse
Predicted Molecular Mass	17.7 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Mouse OX40L (N-8His)
Catalog #: PHM1742



Alternative Names

Tumor necrosis factor ligand superfamily member 4; OX40 ligand; OX40L; CD252; Tnfsf4

Background

OX40 ligand (OX40L), also called CD252, is a single-pass type II membrane protein of the TNF/TNF receptor superfamily. OX40L is expressed by DCs, macrophages and B cells and signals via its cognate receptor OX40 which is mainly expressed on APCs. OX40L/OX40 interactions are important in T-cell activation and survival and for the generation of memory T cells from activated effector T cells. OX40L–OX40 co-stimulation leads to activation of TNF receptor associated factor (TRAF) 2, 3 and 5. This pathway has been shown to prolong the survival of effector CD4⁺Th cells as well as contributes to generation of memory T cells.

Note

For Research Use Only , Not for Diagnostic Use.