


Summary

Name	CD38/ADP-ribosyl Cyclase 1/cyclic ADP-ribose Hydrolase 1
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/µg as determined by LAL test.
Construction	Recombinant Mouse ADP-ribosyl Cyclase/cyclic ADP-ribose Hydrolase 1 is produced by our Mammalian expression system and the target gene encoding Leu45-Thr304 is expressed with a 6His tag at the C-terminus.
Accession #	P56528
Host	Human Cells
Species	Mouse
Predicted Molecular Mass	30.6 KDa
Formulation	Supplied as a 0.2 μ m filtered solution of 20mM Tris-HCl, 150mM NaCl, pH 8.0.
Shipping	The product is shipped on dry ice/polar packs. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	

SDS-PAGE image



Background

Alternative Names	ADP-ribosyl cyclase/cyclic ADP-ribose hydrolase 1; 2-phospho-ADP-ribosyl cyclase; 2-phospho-cyclic-ADP-ribose transferase; ADP-ribosyl cyclase 1; Cyclic ADP-ribose
Background	hydrolase 1; cADPr hydrolase 1; T10; CD38 CD38, also known as ADP-ribosyl cyclase/cyclic ADP-ribose hydrolase 1, is a Signal-
Background	CD38, also known as ADP-ribosyl cyclase/cyclic ADP-ribose hydrolase 1, is a Sign



anchor for type II membrane protein. CD38 is able to transform NAD+ to ADP-Dribose and nicotinamide. It also can transform NADP+ to nicotinate-adenine dinucleotide phosphate and nicotinamide. CD38 is expressed at high levels in pancreas, liver, kidney, brain, testis, ovary, placenta, malignant lymphoma and neuroblastoma. Synthesizes the second messagers cyclic ADP-ribose and nicotinate-adenine dinucleotide phosphate, the former a second messenger for glucose-induced insulin secretion. Also has cADPr hydrolase activity. Also moonlights as a receptor in cells of the immune system.

Note

For Research Use Only, Not for Diagnostic Use.