


Product Name: Recombinant Mouse TROP-2 (C-6His)
Catalog #: PHM1926



Summary

Name	TROP-2/TACSTD2/Tumor-associated Calcium Signal Transducer 2
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Mouse Tumor-associated Calcium Signal Transducer 2 is produced by our Mammalian expression system and the target gene encoding Gln25-Gly270 is expressed with a 6His tag at the C-terminus.
Accession #	Q8BGV3
Host	Human Cells
Species	Mouse
Predicted Molecular Mass	28.8 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Mouse TROP-2 (C-6His)
Catalog #: PHM1926



Alternative Names

Tumor-associated calcium signal transducer 2; Tacstd2; Trop2; Cell surface glycoprotein Trop-2

Background

Tumor-associated calcium signal transducer 2(Tacstd2), also known as Cell surface glycoprotein Trop-2, belongs to the EPCAM family. Tacstd2 expressed in kidney, lung, ovary and testis and has high levels of expression in immortalized keratinocytes. Tacstd2 may functions as a growth factor receptor. It has negative regulation of branching involved in ureteric bud morphogenesis, cell motility, epithelial cell migration, ruffle assembly, stress fiber assembly and substrate adhesion-dependent cell spreading. Also, it has positive regulation of stem cell proliferation. Tacstd2 is capable of transducing an intracellular calcium signal and may play a role in tumor growth. It also has adhesive functions.

Note

For Research Use Only , Not for Diagnostic Use.