

Product Name: Recombinant Mouse Siglec-15 (C-6His)
Catalog #: PHM2075




Summary

Name	Siglec-15/CD33L3/Sialic acid-binding Ig-like lectin 15/CD33 antigen-like 3
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Mouse Sialic Acid-binding Ig-like Lectin 15 is produced by our Mammalian expression system and the target gene encoding Arg24-Thr262 is expressed with a 6His tag at the C-terminus.
Accession #	A7E1W8
Host	Human Cells
Species	Mouse
Predicted Molecular Mass	26.5 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, 150mM NaCl, 5% Thehalose, 0.3% Chaps, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Product Name: Recombinant Mouse Siglec-15 (C-6His)
Catalog #: PHM2075



Alternative Names

Sialic acid-binding Ig-like lectin 15; Siglec-15; CD33 antigen-like 3; CD33L3

Background

Human Siglec-15 is a transmembrane glycoprotein in the Siglec family. Siglecs are type I transmembrane proteins where the NH₃⁺-terminus is in the extracellular space and the COO⁻-terminus is cytosolic. Each Siglec contains an N-terminal V-type immunoglobulin domain (Ig domain) which acts as the binding receptor for sialic acid. These lectins are placed into the group of I-type lectins because the lectin domain is an immunoglobulin fold. All Siglecs are extended from the cell surface by C2-type Ig domains which have no binding activity. Siglecs differ in the number of these C2-type domains. Human Siglec-15 consists of a 244 amino acid (aa) extracellular domain (ECD) with two Ig-like domains, a 21 aa transmembrane segment, and a 44 aa cytoplasmic domain. Siglec-15 function is important for osteoclast formation and TRANCE/RANK Ligand signaling in osteoclasts

Note

For Research Use Only , Not for Diagnostic Use.