Product Name: Recombinant Mouse CD79B (C-Fc)

Catalog #: PHM2114



Summary

Name CD79B/B29

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/µg as determined by LAL test.

Construction Recombinant Mouse CD79B is produced by our Mammalian expression

system and the target gene encoding Val26-Asp158 is expressed with a

human IgG1 Fc tag at the C-terminus.

Accession # P15530

Host **Human Cells**

Species Mouse

Predicted Molecular Mass 42 KDa

Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. **Formulation**

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 Stability&Storage


months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Mouse CD79B (C-Fc) Catalog #: PHM2114



Alternative Names B-Cell Antigen Receptor Complex-Associated Protein Beta Chain; B-Cell-Specific

Glycoprotein B29; Ig-Beta; Immunoglobulin-Associated B29 Protein; CD79b;

CD79B; B29; IGB

Background CD79B is a single-pass type I membrane protein. CD79B contains one Ig-like V-

type domain and one ITAM domain. CD79B is required in cooperation with CD79A for initiation of the signal transduction cascade activated by the B-cell antigen receptor complex (BCR), which leads to internalization of the complex, trafficking to late endosomes and antigen presentation. CD79B enhances phosphorylation of CD79A, possibly by recruiting kinases that phosphorylate CD79A or by recruiting

proteins that bind to CD79A and protect it from dephosphorylation.

Note

For Research Use Only, Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838