## **Product Name: Recombinant Mouse IL-1RL2 (C-6His)** Catalog #: PHM2131



### **Summary**

Name IL-36R/IL-1RL2/IL-1Rrp2/IL-1 Receptor-Like 2

**Purity** Greater than 95% as determined by reducing SDS-PAGE

**Endotoxin level** <1 EU/µg as determined by LAL test.

Construction Recombinant Mouse Interleukin-1 Receptor-like 2 is produced by our

Mammalian expression system and the target gene encoding Asp22-

Arg338(Ala158Val) is expressed with a 6His tag at the C-terminus.

Accession # Q9ERS7

Host **Human Cells** 

**Species** Mouse

**Predicted Molecular Mass** 36.9 KDa

Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. **Formulation** 

**Shipping** The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 Stability&Storage


months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

#### **SDS-PAGE** image



## **Background**

# Product Name: Recombinant Mouse IL-1RL2 (C-6His) Catalog #: PHM2131



Alternative Names Interleukin-1 receptor-like 2; IL1RL2; IL-36 receptor; IL-36R; IL-1Rrp2; IL1R-rp2

**Background** Interleukin-1 receptor-like 2 is a protein that in humans is encoded by the IL1RL2

gene, belongs to the interleukin-1 receptor family.IL1RL2 is the receptor for interleukin-36 (IL36A, IL36B and IL36G). After binding to interleukin-36 associates with the coreceptor IL1RAP to form the interleukin-36 receptor complex which mediates interleukin-36-dependent activation of NF-kappa-B, MAPK and other

pathways.

#### Note

For Research Use Only, Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838