# **Product Name: Recombinant Mouse CD27L (N-10His)**

Catalog #: PHM2243



### **Summary**

Name CD27L/CD70/CD27 Ligand/TNFSF7

**Purity** Greater than 95% as determined by reducing SDS-PAGE

**Endotoxin level** <1 EU/μg as determined by LAL test.

Construction Recombinant Mouse CD27 Ligand/CD70 is produced by our Mammalian

expression system and the target gene encoding Gln47-Pro195 is expressed

with a 10His tag at the N-terminus.

Accession # Q05A52

**Host** Human Cells

**Species** Mouse

Predicted Molecular Mass 19 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

**Stability&Storage** Store at  $\leq$ -70°C, stable for 6 months after receipt. Store at  $\leq$ -70°C, stable for 3

months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

**Reconstitution** Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.


### **SDS-PAGE** image

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

# **Product Name: Recombinant Mouse CD27L (N-10His)**

Catalog #: PHM2243





#### **Alternative Names**

CD70 antigen; CD70; CD27 ligand; Tumor necrosis factor ligand superfamily member 7; CD27LG; TNFSF7; CD27L

## **Background**

CD27 Ligand (CD70) is a 30 kDa type 2 transmembrane glycoproteinis that belongs to the tumor necrosis factor (TNF) ligand family. Human CD27 Ligand extracellular region is 64% aa identical to mouse and rat CD27 Ligand extracellular regions. This cytokine is a ligand for TNFRSF27/CD27. It is a surface antigen on activated, but not on resting, T and B lymphocytes. It induces proliferation of costimulated T cells, enhances the generation of cytolytic T cells, and contributes to T cell activation. This cytokine is also reported to play a role in regulating B-cell activation, cytotoxic function of natural killer cells, and immunoglobulin sythesis.

#### Note

For Research Use Only, Not for Diagnostic Use.