


Product Name: Recombinant Mouse TNF RII (C-Fc)
Catalog #: PHM2277



Summary

Name	TNF RII/TNFRSF1B/CD120b/TNFR2/TNF Receptor II/Tumor Necrosis Factor Receptor II
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Mouse Tumor Necrosis Factor Receptor II is produced by our Mammalian expression system and the target gene encoding Val23-Gly258 is expressed with a human IgG1 Fc tag at the C-terminus.
Accession #	Q545P4
Host	Human Cells
Species	Mouse
Predicted Molecular Mass	52.3 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Mouse TNF RII (C-Fc)
Catalog #: PHM2277



Alternative Names

Tumor necrosis factor receptor superfamily member 1b; Tnfrsf1b

Background

Tumor Necrosis Factor Receptor Superfamily Member 1B (TNFRSF1B) is a member of the Tumor Necrosis Factor Receptor Superfamily. TNFRSF1B contains four TNFR-Cys repeats. TNFRSF1B can be cleaved into the following 2 chains: Tumor necrosis factor receptor superfamily member 1b and membrane form and Tumor necrosis factor-binding protein 2. TNFRSF1B is a receptor with high affinity for TNFSF2/TNF- α and approximately 5-fold lower affinity for homotrimeric TNFSF1/lymphotoxin- α . TNFRSF1B mediates most of the metabolic effects of TNF- α . TNF- α -induced apoptosis suggests that it regulates TNF- α function by antagonizing its biological activity.

Note

For Research Use Only , Not for Diagnostic Use.