Product Name: Recombinant Rat IL-2 (C-6His)

Catalog #: PHR0994



Summary

Name IL-2/Interleukin-2/T cell growth factor/TCGF

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/μg as determined by LAL test.

Construction Recombinant Rat Interleukin-2 is produced by our Mammalian expression

system and the target gene encoding Ala21-Gln155 is expressed with a 6His

tag at the C-terminus.

Accession # P17108

Host Human Cells

Species Rat

Predicted Molecular Mass 16.3 KDa

Formulation Supplied as a 0.2 µm filtered solution of 5mM NaH2POsub//u003E4/sub//u003E,

5mM Citric acid, 150mM NaCl, pH 4.0.

Shipping The product is shipped on dry ice/polar packs. Upon receipt, store it immediately

at the temperature listed below.

Stability&Storage Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3

months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

Reconstitution 0.00.0


SDS-PAGE image

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Product Name: Recombinant Rat IL-2 (C-6His)

Catalog #: PHR0994





Alternative Names

Interleukin-2; IL-2; T-cell growth factor; TCGF; Aldesleukin; IL2

Background

Interleukin-2(IL-2)is a O-glycosylated four α -helix bundle cytokine that has potent stimulatory activity for antigenactivated T cells. It is expressed by CD4+ and CD8+ T cells, $\gamma\delta$ T cells, B cells, dendritic cells, and eosinophils. Mature rat IL-2 shares 66% and 73% amino acid sequence identity with human and mouse IL-2,respectively. The receptor for IL-2 consists of three subunits that are present on the cell surface in varying preformed complexes. IL-2 is a powerful immunoregulatory lymphokine produced by T-cells in response to antigenic or mitogenic stimulation. IL-2/IL-2R signaling is required for T-cell proliferation and other fundamental functions that are essential for the immune response. IL-2 stimulates growth and differentiation of B-cells, NK cells, lymphokine-activated killer cells, monocytes, macrophages and oligodendrocytes.

Note

For Research Use Only, Not for Diagnostic Use.