

Product Name: Recombinant Cynomolgus TIM-3 (C-Fc)
Catalog #: PHV1652




Summary

Name	TIM-3/HAVCR2/TIMD3/T Cell Immunoglobulin and Mucin Domain-3
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Cynomolgus T Cell Immunoglobulin And Mucin Domain-3 is produced by our Mammalian expression system and the target gene encoding Ser22-Arg201 is expressed with a human IgG1 Fc tag at the C-terminus.
Accession #	G7P6Q7
Host	Human Cells
Species	Cynomolgus
Predicted Molecular Mass	46.3 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Product Name: Recombinant Cynomolgus TIM-3 (C-Fc)
Catalog #: PHV1652



Alternative Names

T cell immunoglobulin and mucin domain3; HAVCR2; Tim-3; TIM3

Background

T cell immunoglobulin and mucin domain 3 is a member of the TIM family of immune regulating molecules. Mature cynomolgus TIM3 consists of a 182 amino acid (aa) extracellular domain (ECD), a 21 aa transmembrane segment, and a 78 aa cytoplasmic tail. TIM3 is up-regulated on several populations of activated myeloid cells (macrophage, monocyte, dendritic cell, microglia, mast cell) and T cells (Th1, CD8+, NK, Treg). Its binding to Galectin9 induces a range of immunosuppressive functions which enhance immune tolerance and inhibit anti-tumor immunity. TIM3 ligation attenuates CD8+ and Th1 cell responses and promotes the activity of Treg and myeloid derived suppressor cells. TIM3 interactions with Galectin-9 can trigger immune stimulatory effects, such as the coactivation of NK cell cytotoxicity.

Note

For Research Use Only , Not for Diagnostic Use.