


Product Name: Recombinant Cynomolgus CD3d (C-Fc)
Catalog #: PHV1939



Summary

Name	CD3D/CD3 delta/T-cell surface glycoprotein CD3 delta chain
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Cynomolgus T-cell Surface Glycoprotein CD3 Delta Chain is produced by our Mammalian expression system and the target gene encoding Phe22-Ala105 is expressed with a human IgG1 Fc tag at the C-terminus.
Accession #	Q95LI8
Host	Human Cells
Species	Cynomolgus
Predicted Molecular Mass	36.7 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Product Name: Recombinant Cynomolgus CD3d (C-Fc)
Catalog #: PHV1939



Background

Alternative Names

T-cell surface glycoprotein CD3 delta chain; T-cell receptor T3 delta chain; CD3d; CD3D

Background

T-cell surface glycoprotein CD3 delta chain (CD3D) is a single-pass type I membrane protein. CD3D, together with CD3-gamma, CD3-epsilon and CD3-zeta, and the T-cell receptor alpha/beta and gamma/delta heterodimers, forms the T cell receptor-CD3 complex. CD3 chains are present as CD3gammaepsilon, deltaepsilon, and zeta zeta dimers in the receptor complex and play critical roles in the antigen receptor assembly, transport to the cell surface, and the receptor-mediated signal transduction. T cell receptor-CD3 complex plays an important role in coupling antigen recognition to several intracellular signal-transduction pathways. This complex is critical for T-cell development and function, and represents one of the most complex transmembrane receptors. The T cell receptor-CD3 complex is unique in having ten cytoplasmic immunoreceptor tyrosine-based activation motifs (ITAMs). CD3D contains 1 ITAM domain and has been shown to interact with CD8A.

Note

For Research Use Only , Not for Diagnostic Use.