

Product Name: Recombinant Cynomolgus 5T4 (C-6His)
Catalog #: PHV2128




Summary

Name	TPBG/5T4/Trophoblast glycoprotein
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Cynomolgus Trophoblast Glycoprotein is produced by our Mammalian expression system and the target gene encoding Thr35-Ser355 is expressed with a 6His tag at the C-terminus.
Accession #	Q4R8Y9
Host	Human Cells
Species	Cynomolgus
Predicted Molecular Mass	35.7 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Product Name: Recombinant Cynomolgus 5T4 (C-6His)
Catalog #: PHV2128



Alternative Names

TPBG; 5T4; 5T4AG; M6P1; WAIF1

Background

Trophoblast glycoprotein, also known as TPBG and 5T4, it belongs to the LRR family of proteins. The C-terminus of LRR N-terminal cap (LRRNT) and LRR 1 are essential for the inhibition of the Wnt signaling pathway. TPBG may function as an inhibitor of Wnt/beta-catenin signaling by indirectly interacting with LRP6 and blocking Wnt3a-dependent LRP6 internalization. It expressed by all types of trophoblasts as early as 9 weeks of development. In adult tissues, the expression is limited to a few epithelial cell types but is found on a variety of carcinoma.

Note

For Research Use Only , Not for Diagnostic Use.