Catalog #: PHV2272



Summary

ACE-2/Angiotensin-Converting Enzyme 2 Name

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/µg as determined by LAL test.

Construction Recombinant Rhesus Macaque Angiotensin-Converting Enzyme 2 is produced

by our Mammalian expression system and the target gene encoding Gln18-

Val739 is expressed with a 10His tag at the C-terminus.

Accession # ACI04564.1

Host **Human Cells**

Species Rhesus Macaque

Predicted Molecular Mass 85.1 KDa

Formulation Supplied as a 0.2 µm filtered solution of 25mM Tris-HCl, 150mM NaCl, 1mM

ZnCl2, pH 7.5.

The product is shipped on dry ice/polar packs. Upon receipt, store it immediately **Shipping**

at the temperature listed below.

Stability&Storage Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3

months under sterile conditions after opening. Please minimize freeze-thaw


cycles.

Reconstitution

SDS-PAGE image

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838





Alternative Names

Angiotensin-Converting Enzyme 2; ACE-Related Carboxypeptidase; Angiotensin-Converting Enzyme Homolog; ACEH; Metalloprotease MPROT15; ACE2

Background

Angiotensin-Converting Enzyme 2 (ACE-2) is an integral membrane protein and a zinc metalloprotease of the ACE family, the ACE family includes somatic and germinal ACE. ACE-2 cleaves angiotensins I and II as a carboxypeptidase, ACE-2 converts angiotensin I to angiotensin 1-9, and angiotensin II to angiotensin 1-7. ACE-2 is also able to hydrolyze apelin-13 and dynorphin-13 with high efficiency. ACE-2 can be high expressed in testis, kidney and heart, in colon, small intestine and ovary at moderate levels. Captopril and lisinopril as the classical ACE inhibitor don't inhibit ACE-2 activity. ACE-2 may play an important role in regulating the heart function.

Note

For Research Use Only, Not for Diagnostic Use.