Catalog #: PHV2276



Summary

TROP-2/TACSTD2/Tumor-associated Calcium Signal Transducer 2 Name

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/µg as determined by LAL test.

Construction Recombinant Rhesus Macaque Tumor-associated Calcium Signal Transducer

2 is produced by our Mammalian expression system and the target gene

encoding His27-Arg272 is expressed with a 6His tag at the C-terminus.

Accession # XP 001114599.1

Host **Human Cells**

Species Rhesus Macaque

Predicted Molecular Mass 28.7 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

The product is shipped at ambient temperature. Upon receipt, store it **Shipping**

immediately at the temperature listed below.

Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 Stability&Storage


months under sterile conditions after opening. Please minimize freeze-thaw

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

> not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image





Alternative Names

Tumor-associated calcium signal transducer 2; Membrane component chromosome 1 surface marker 1; Cell surface glycoprotein Trop-2; TACSTD2; TROP2

Background

Tumor associated calcium signal transducer 2 (TACSTD2, TROP-2) is a type I cell surface glycoprotein that is highly expressed on human carcinomas. It was originally identified as an antigen present on human gastrointestinal tumors and is the second of two members of this family. Human and mouse TROP-2 share 87% amino acid (aa) similarity. TROP-2 is capable of transducing an intracellular calcium signal and may play a role in tumor growth. It also has adhesive functions.

Note

For Research Use Only, Not for Diagnostic Use.