# Product Name: Recombinant Cynomolgus Neuropilin-1 (C-6His) Enkille

## **Summary**

Name Neuropilin-1/NRP1

**Purity** Greater than 95% as determined by reducing SDS-PAGE

**Endotoxin level** <1 EU/μg as determined by LAL test.

Construction Recombinant Cynomolgus Neuropilin-1 is produced by our Mammalian

expression system and the target gene encoding Phe22-Gly613 is expressed

with a 6His tag at the C-terminus.

Accession # G7PEQ1

**Host** Human Cells

**Species** Cynomolgus

Predicted Molecular Mass 67.5 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

**Stability&Storage** Store at  $\leq$ -70°C, stable for 6 months after receipt. Store at  $\leq$ -70°C, stable for 3

months under sterile conditions after opening. Please minimize freeze-thaw


cycles.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

### **SDS-PAGE** image

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838



#### **Alternative Names**

CD304; NRP1; NRPNP1; VEGF165R; BDCA4

# **Background**

Neuropilin-1 (Npn-1, previously neuropilin; also CD304) is a 130 - 140 kDa type I transmembrane (TM) glycoprotein that regulates axon quidance and angiogenesis. Two homologues, Neuropilin-1 and Neuropilin-2, are identified. Neuropilin-1 binds to semaphorin 3A, The PLGF-2 isoform of PGF, The VEGF-165 isoform of VEGF and VEGF-B. Coexpression with KDR results in increased VEGF-165 binding to KDR as well as increased chemotaxis. It may regulate VEGF-induced angiogenesis. The soluble isoform 2 binds VEGF-165 and appears to inhibit its binding to cells. NRP1 expression is regulated in EC by tumor necrosis factoralpha, the transcription factors dHAND and Ets-1, and vascular injury. NRP1 upregulation is positively correlated with the progression of various tumors.

#### Note

For Research Use Only, Not for Diagnostic Use.