

Product Name: Recombinant Cynomolgus TRAIL R2 (C-6His)
Catalog #: PHV2408




Summary

Name	TRAIL R2/TNFRSF10B/CD262/DR5/KILLER/Fas-like protein
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Cynomolgus TNF-Related Apoptosis-Inducing Ligand Receptor 2 is produced by our Mammalian expression system and the target gene encoding Ile58-Ser212 is expressed with a 6His tag at the C-terminus.
Accession #	A0A2K5TXK0
Host	Human cells
Species	Cynomolgus
Predicted Molecular Mass	18 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Product Name: Recombinant Cynomolgus TRAIL R2 (C-6His)
Catalog #: PHV2408



Alternative Names

Tumor Necrosis Factor Receptor Superfamily Member 10B; Death Receptor 5; TNF-Related Apoptosis-Inducing Ligand Receptor 2; TRAIL Receptor 2; TRAIL-R2; CD262; TNFRSF10B; DR5; KILLER; TRAILR2; TRICK2; ZTNFR9

Background

TNFRSF10B is a member of the TNF-receptor superfamily, and contains an intracellular death domain. This receptor can be activated by tumor necrosis factor-related apoptosis inducing ligand (TNFSF10/TRAIL/APO-2L), and transduces apoptosis signal. The adapter molecule FADD recruits caspase-8 to the activated receptor and is required for the apoptosis mediated by TNFRSF10B. TNFRSF10B is expressed in a number of cell types, and to particularly high levels in lymphocytes and spleen. This single-pass transmembrane protein contains two cysteine-rich repeat units in its extracellular region, followed by a transmembrane segment and a cytoplasmic tail containing a typical "death domain". TNFRSF10B expression is regulated by the tumor suppressor p53. It is also indicated that the activation of NF-kappa-B can be promoted by TNFRSF10B.

Note

For Research Use Only , Not for Diagnostic Use.