


Summary

Name	VEGF164/VEGFA/Vascular Endothelial Growth Factor Isoform 164
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<0.1 EU/μg as determined by LAL test.
Construction	Recombinant Rat Vascular Endothelial Growth Factor A is produced by our Yeast expression system and the target gene encoding Ala27-Arg190(Ala36Thr) is expressed.
Accession #	P16612-2
Host	P. pastoris
Species	Rat
Predicted Molecular Mass	19.2 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Product Name: Recombinant Rat VEGF 164
Catalog #: PPR1807



Alternative Names

Vascular endothelial growth factor A; Vascular permeability factor; VEGF; VEGF-A; VPF

Background

Vascular endothelial growth factor (VEGF/VEGF-A) is originally known as vascular permeability factor (VPF). It belongs to the PDGF family with a cysteine-knot structure comprised of eight conserved cysteine residues, and reckoned as a potent mediator in the process of angiogenesis and vasculogenesis in either fetus or adult. VEGF is particularly expressed in supraoptic , paraventricular nuclei and the choroid plexus of the pituitary, and abundant in the corpus luteum of the ovary and in kidney glomeruli. The rat VEGF protein contains a putative 20 amino acids (aa) signal peptide, and alternative splicing of rat VEGF gene produces isoforms of 120, 144, 164 and 188 aa. Rat VEGF164 respectively displays 97% and 88% aa identity with that regions of mouse and human VEGF. VEGF can bind to the FLT1/VEGFR1 and KDR/VEGFR2 receptors, heparan sulfate and heparin, and play important roles in inducing endothelial cell proliferation, promoting cell migration, inhibiting apoptosis and inducing permeabilization of blood vessels.

Note

For Research Use Only , Not for Diagnostic Use.