Product Name: CD10 (7C7) Mouse Monoclonal Antibody Enkilife Catalog #: AMM00733

Summary

Production Name CD10 (7C7) Mouse Monoclonal Antibody

Description Mouse Monoclonal Antibody

Host Mouse Application IHC-P

Reactivity Human, Rat, Mouse

Performance

ConjugationUnconjugatedModificationUnmodified

Isotype IgG1

Clonality Monoclonal Form Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw Storage

cycles.

Buffer Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.

Purification Affinity Purification

Immunogen

Gene Name MME

MME; EPN; Neprilysin; Atriopeptidase; Common acute lymphocytic leukemia antigen;

Alternative Names CALLA; Enkephalinase; Neutral endopeptidase 24.11; NEP; Neutral endopeptidase; Skin

fibroblast elastase; SFE; CD10

 Gene ID
 4311

 SwissProt ID
 P08473.

Application

Dilution Ratio IHC: 1:50-1:100

Molecular Weight -




Background

CD10 is a transmembrane type II molecule and functions as a metallo-peptidase requiring zinc. Specifically, CD10 cleaves 1-3 amino-terminal amino acids from peptides with a preference for neutral amino acids (valine, iso-leucine, phenylalanine, leucine or alanine). Involved in the degradation of atrial natriuretic factor (ANF). Displays UV-inducible elastase activity toward skin preelastic and elastic fibers.


Research Area

Immunology

Image Data



Immunofluorescence analysis of CD10 (7C7) in Human uterus tissue using CD10 (7C7) antibody(red), and DAPI (blue).




Immunofluorescence analysis of CD10 (7C7) in rat kidney using CD10 antibody(red), and DAPI (blue).


Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Product Name: CD10 (7C7) Mouse Monoclonal Antibody Enkilife Catalog #: AMM00733






Immunohistochemistry analysis of paraffin-embedded Human uterus tissue using CD10 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval. Negative control was used by secondary antibody only.




Immunohistochemistry analysis of paraffin-embedded rat kidney tissue using CD10 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval. Negative control was used by secondary antibody only.



Immunohistochemistry analysis of paraffin-embedded mouse lung tissue using CD10 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval. Negative control was used by secondary antibody only.





Immunohistochemistry analysis of paraffin-embedded Human breast cancer using CD10 (7C7) antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.v

Note

For research use only.