

**Product Name: MAP2 (3B5) Mouse Monoclonal Antibody**  
**Catalog #: AMM03341**

---

## Summary

<b>Production Name</b>	MAP2 (3B5) Mouse Monoclonal Antibody
<b>Description</b>	Mouse Monoclonal Antibody
<b>Host</b>	Mouse
<b>Application</b>	IHC-P, ICC/IF
<b>Reactivity</b>	Human, Mouse, Rat

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG1
<b>Clonality</b>	Monoclonal
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.
<b>Purification</b>	Affinity Purification

## Immunogen

<b>Gene Name</b>	MAP2
<b>Alternative Names</b>	Microtubule associated protein 2; MAP2A; MAP2B; MAP2C
<b>Gene ID</b>	4133
<b>SwissProt ID</b>	P11137.

## Application

<b>Dilution Ratio</b>	IHC: 1:50-1:100 IF: 1:50-1:200
<b>Molecular Weight</b>	-

## Background

---

**Product Name: MAP2 (3B5) Mouse Monoclonal Antibody**  
**Catalog #: AMM03341**

---




The exact function of MAP2 is unknown but MAPs may stabilize the microtubules against depolymerization. They also seem to have a stiffening effect on microtubules.


## Research Area

Neuroscience

## Image Data




Immunohistochemistry analysis of paraffin-embedded Human Tonsil tissue using MAP2 (3B5) antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval. Negative control was used by secondary antibody only.




Immunohistochemical analysis of paraffin-embedded Human tonsils using MAP2 (3B5) antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval. Negative control was used by secondary antibody only.

**Product Name: MAP2 (3B5) Mouse Monoclonal Antibody**  
**Catalog #: AMM03341**


---



Immunohistochemistry analysis of paraffin-embedded mouse brain tissue using MAP2 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval. Negative control was used by secondary antibody only.




Immunofluorescence analysis of MAP2 (3B5) in mouse brain tissue using MAP2 (3B5) antibody(7D4)(red),and DAPI (blue).



Immunofluorescence analysis of MAP2 in rat brain using MAP2 antibody(7D4)(red) ,and DAPI (blue).

**Product Name: MAP2 (3B5) Mouse Monoclonal Antibody**  
**Catalog #: AMM03341**

---



Immunohistochemistry analysis of paraffin-embedded Human brain tissue using MAP2 (3B5) antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

### **Note**

For research use only.