Product Name: eNOS(Mix)Mouse Monoclonal Antibody Catalog #: AMM10478

Summary

Production Name eNOS(Mix)Mouse Monoclonal Antibody

Description Mouse Monoclonal Antibody

Host Mouse
Application WB

Reactivity Human, Mouse, Rat, Rabbit

Performance

ConjugationUnconjugatedModificationUnmodified

Isotype IgG

Clonality Monoclonal Form Liquid

Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

PBS, pH 7.4, containing 0.5%BSA, 0.02% New type preservative N as Preservative and

50% Glycerol.

Purification Affinity purification

Immunogen

Buffer

Gene Name NOS3

NOS3; Nitric oxide synthase, endothelial; Constitutive NOS; cNOS; EC-NOS; Endothelial

NOS; eNOS; NOS type III; NOSIII

Gene ID 4846.0

SwissProt ID P29474. Recombinant Protein of eNOS

Application

Dilution Ratio WB 1:500-2000

Molecular Weight 130-140kDa


Background

Nitric oxide is a reactive free radical which acts as a biologic mediator in several processes, including neurotransmission and antimicrobial and antitumoral activities. Nitric oxide is synthesized from L-arginine by nitric oxide synthases. Variations in this gene are associated with susceptibility to coronary spasm. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, May 2009],catalytic activity:L-arginine + n NADPH + n H(+) + m O(2) = citrulline + nitric oxide + n NADP(+),,cofactor:Binds 1 FAD.,cofactor:Binds 1 FMN.,cofactor:Heme group.,cofactor:Tetrahydrobiopterin (BH4). May stabilize the dimeric form of the enzyme.,enzyme regulation:Stimulated by calcium/calmodulin. Inhibited by NOSIP and NOSTRIN.,function:Produces nitric oxide (NO) which is implicated in vascular smooth muscle relaxation through a cGMP-mediated signal transduction pathway. NO mediates vascular endothelial growth factor (VEGF)-induced angiogenesis in coronary vessels and promotes blood clotting through the activation of platelets.,online information:Nitric oxide synthase entry,polymorphism:Variation in NOS3 seem to be associated with susceptibility to coronary spasm.,similarity:Belongs to the NOS family.,similarity:Contains 1 FAD-binding FR-type domain.,similarity:Contains 1 flavodoxin-like domain.,subcellular location:Specifically associates with actin cytoskeleton in the G2 phase of the cell cycle; which is favored by interaction with NOSIP and results in a reduced enzymatic activity, subunit:Homodimer. Interacts with NOSIP and NOSTRIN.,tissue specificity:Platelets, placenta, liver and kidney.,

Research Area

Arginine and proline metabolism; Calcium; VEGF;

Image Data



Western blot analysis of Rat Heart Tissue, diluted at 1:1000.

Note

For research use only.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838