**Antibody** 

Catalog #: AMM16407



## Summary

**Production Name** PPAR Delta(2F9)Mouse Monoclonal Antibody

**Description** Mouse Monoclonal Antibody

**Host** Mouse

**Application** IF-P,IF-F,ICC/IF,IHC-P **Reactivity** Human,Rat,Mouse

#### **Performance**

ConjugationUnconjugatedModificationUnmodified

**Isotype** IgG

**Clonality** Monoclonal

Form Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw  $\bf Storage$ 

cycles.

**Buffer** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.

**Purification** Affinity purification

### **Immunogen**

Gene Name PPARD

Peroxisome proliferator-activated receptor delta (PPAR-delta) (NUCI) (Nuclear

Alternative Names hormone receptor 1) (NUC1) (Nuclear receptor subfamily 1 group C member 2)

(Peroxisome proliferator-activated receptor beta) (PPAR-beta)

**Gene ID** 5467.0

SwissProt ID O03181.Recombinant Protein of PPAR Delta of PPAR Delta

## **Application**

**Dilution Ratio** IF-P/IF-F/ICC/IF 1:50-200, IHC-P 1:100-200

Molecular Weight 50kDa

**Antibody** 

Catalog #: AMM16407




### **Background**

peroxisome proliferator activated receptor delta(PPARD) Homo sapiens This gene encodes a member of the peroxisome proliferator-activated receptor (PPAR) family. PPARs are nuclear hormone receptors that bind peroxisome proliferators and control the size and number of peroxisomes produced by cells. PPARs mediate a variety of biological processes, and may be involved in the development of several chronic diseases, including diabetes, obesity, atherosclerosis, and cancer. This protein is a potent inhibitor of ligand-induced transcription activity of PPAR alpha and PPAR gamma. It may function as an integrator of transcription repression and nuclear receptor signaling. The expression of this gene is found to be elevated in colorectal cancer cells. The elevated expression can be repressed by adenomatosis polyposis coli (APC), a tumor suppressor protein related to APC/beta-catenin signaling pathway. Knockout studies in mice suggested the role of thisfunction:Receptor that binds peroxisome proliferators such as hypolipidemic drugs and fatty acids. Once activated by a ligand, the receptor binds to a promoter element in the gene for acyl-CoA oxidase and activates its transcription. It therefore controls the peroxisomal beta-oxidation pathway of fatty acids. Decreases expression of NPC1L1 once activated by a ligand, online information:Peroxisome proliferator-activated receptor entry, similarity:Belongs to the nuclear hormone receptor family. NR1 subfamily, similarity:Contains 1 nuclear receptor DNA-binding domain., subunit:Heterodimer with the retinoid X receptor, tissue specificity:Ubiquitous with maximal levels in placenta and skeletal muscle.

#### **Research Area**

PPAR;WNT;WNT-T CELLPathways in cancer;Acute myeloid leukemia;

#### **Image Data**




Immunofluorescence analysis of rat-spleen tissue. 1,PPAR Delta Mouse Monoclonal Antibody (2F9) (red) was diluted at 1:200 (4°C,overnight) . 2, Cy3 labled Secondary antibody was diluted at 1:300 (room temperature, 50min) .3, Picture B: DAPI (blue) 10min. Picture A:Target. Picture B: DAPI. Picture C: merge of A+B

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838


**Antibody** 

Catalog #: AMM16407






Immunofluorescence analysis of mouse-spleen tissue. 1,PPAR Delta Mouse Monoclonal Antibody (2F9) (red) was diluted at 1:200 (4°C,overnight) . 2, Cy3 labled Secondary antibody was diluted at 1:300 (room temperature, 50min) .3, Picture B: DAPI (blue) 10min. Picture A:Target. Picture B: DAPI. Picture C: merge of A+B



Immunohistochemical analysis of paraffin-embedded Human-uterus tissue. 1,PPAR Delta Mouse Monoclonal Antibody (2F9) was diluted at 1:200 (4°C,overnight) . 2, Sodium citrate pH 6.0 was used for antibody retrieval (>98°C,20min) . 3,Secondary antibody was diluted at 1:200 (room tempeRature, 30min) . Negative control was used by secondary antibody only.



Immunohistochemical analysis of paraffin-embedded Human Brain Tissue using PPAR Delta Mouse mAb diluted at 1:200.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Antibody

Catalog #: AMM16407





Immunohistochemical analysis of paraffin-embedded Mouse Brain Tissue using PPAR Delta Mouse mAb diluted at 1:200.

#### Note

For research use only.