Product Name: STAT3(3F4)Mouse Monoclonal Antibody Enkilife Catalog #: AMM18356

Summary

Production Name STAT3(3F4)Mouse Monoclonal Antibody

Description Mouse Monoclonal Antibody

Host Mouse

Application WB,IHC-P,IF-P,IF-F,ICC/IF

Reactivity Human, Rat, Mouse

Performance

ConjugationUnconjugatedModificationUnmodified

Isotype IgG

Clonality Monoclonal Form Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw

cycles.

Buffer Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.

Purification Affinity purification

Immunogen

Storage

Gene Name STAT3

STAT3; APRF; Signal transducer and activator of transcription 3; Acute-phase response Alternative Names

factor

Gene ID 6774.0

SwissProt ID P40763.Recombinant Protein of STAT3

Application

Dilution Ratio WB 1:1000-2000, IHC-P 1:100-200, IF-P/IF-F/ICC/IF 1:50-200

Molecular Weight 79,86kDa

Background

Product Name: STAT3(3F4)Mouse Monoclonal Antibody Enkilife Catalog #: AMM18356

The protein encoded by this gene is a member of the STAT protein family. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein is activated through phosphorylation in response to various cytokines and growth factors including IFNs, EGF, IL5, IL6, HGF, LIF and BMP2. This protein mediates the expression of a variety of genes in response to cell stimuli, and thus plays a key role in many cellular processes such as cell growth and apoptosis. The small GTPase Rac1 has been shown to bind and regulate the activity of this protein. PIAS3 protein is a specific inhibitor of this protein. Mutations in this gene are associated with infantile-onset multisystem autoimmune disease and hyperdisease:Defects in STAT3 are the cause of hyperimmunoglobulin E recurrent infection syndrome autosomal dominant (AD-HIES) [MIM:147060]; also known as hyper-lgE syndrome or Job syndrome. AD-HIES is a rare disorder of immunity and connective tissue characterized by immunodeficiency, chronic eczema, recurrent Staphylococcal infections, increased serum IqE, eosinophilia, distinctive coarse facial appearance, abnormal dentition, hyperextensibility of the joints, and bone fractures., function: Transcription factor that binds to the interleukin-6 (IL-6)responsive elements identified in the promoters of various acute-phase protein genes. Activated by IL31 through IL31RA, miscellaneous: Involved in the gp130-mediated signaling pathway, online information: STAT3 entry, online information:STAT3 mutation db,PTM:Tyrosine phosphorylated in response to IL-6, IL-11, CNTF, LIF, CSF-1, EGF, PDGF, IFNalpha and OSM. Phosphorylated on serine upon DNA damage, probably by ATM or ATR. Serine phosphorylation is important for the formation of stable DNA-binding STAT3 homodimers and maximal transcriptional activity., similarity: Belongs to the transcription factor STAT family., similarity: Contains 1 SH2 domain., subcellular location: Shuttles between the nucleus and the cytoplasm. Constitutive nuclear presence is independent of tyrosine phosphorylation., subunit: Forms a homodimer or a heterodimer with a related family member (at least STAT1). Interacts with NCOA1, PELP1, SOCS7 and STATIP1. Interacts with HCV core protein. Interacts with IL23R in presence of IL23. Interacts with IL31RA. Interacts with SIPAR. Interacts (via SH2 domain) with NLK (By similarity). Interacts with KPNA4 and KPNA5; KPNA4 may be the primary mediator of nuclear import (By similarity). Interacts with TMF1, tissue specificity: Heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas.,


Research Area

Chemokine; Jak STAT; Adipocytokine; Pathways in cancer; Pancreatic cancer; Acute myeloid leukemia;


Image Data

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838







Western blot analysis of 1) Rat Heart Tissue, 2) Mouse Heart Tissue, 3) Hela with STAT3 Mouse mAb diluted at 1:2,000.



Immunohistochemical analysis of paraffin-embedded Human Colon Carcinoma using STAT3Mouse mAb diluted at 1:200.



Immunohistochemical analysis of paraffin-embedded Mouse Brain Tissue using STAT3Mouse mAb diluted at 1:200.



Western blot detection of STAT3 in human breast cancer cell line MCF-7 (A), T47D (B), MDA-MB-231 (C) and Cal51 (D) using STAT3 mouse mAb (1:2000 diluted). Predicted band size: 86kDa. Observed band size: 86kDa. Picture was kindly provided by our customer from Tianjin Medical University Cancer Institute and Hospital

Note

For research use only.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838