

**Product Name: cIAP1 Rabbit Monoclonal Antibody**  
**Catalog #: AMRe01826**



## Summary

<b>Production Name</b>	cIAP1 Rabbit Monoclonal Antibody
<b>Description</b>	Rabbit Monoclonal antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB, ICC/IF
<b>Reactivity</b>	Human

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Monoclonal
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA
<b>Purification</b>	Affinity Purification

## Immunogen

<b>Gene Name</b>	BIRC2
<b>Alternative Names</b>	cIAP1; C-IAP1; IAP homolog B; Inhibitor of apoptosis protein 2; IAP-2; hIAP-2; hIAP2; API1; IAP2; MIHB; RNF48
<b>Gene ID</b>	329
<b>SwissProt ID</b>	Q13490.

## Application

<b>Dilution Ratio</b>	WB: 1:500-1:1000 IF: 1:50-1:200
<b>Molecular Weight</b>	Calculated MW: 70 kDa; Observed MW: 70 kDa

**Product Name: cIAP1 Rabbit Monoclonal Antibody**  
**Catalog #: AMRe01826**




## Background

Apoptotic suppressor. The BIR motifs region interacts with TNF receptor associated factors 1 and 2 (TRAF1 and TRAF2) to form an heteromeric complex, which is then recruited to the tumor necrosis factor receptor 2 (TNFR2).


## Research Area

Cell Biology

## Image Data



Western blot analysis of cIAP1 in K562 lysates using cIAP1 antibody.



Immunocytochemistry analysis of cIAP1(green) in HeLa using cIAP1 antibody, and DAPI(blue)

## Note

For research use only.