# Product Name: IKK beta Rabbit Monoclonal Antibody Catalog #: AMRe02148



## **Summary**

Production Name IKK beta Rabbit Monoclonal Antibody

**Description** Rabbit Monoclonal antibody

Host Rabbit
Application WB,IP
Reactivity Human

## **Performance**

ConjugationUnconjugatedModificationUnmodified

**Isotype** IgG

Clonality Monoclonal Form Liquid

**Storage** Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05%

BSA

**Purification** Affinity Purification

### **Immunogen**

Buffer

Gene Name IKBKB

IKBKB; IKKB; Inhibitor of nuclear factor kappa-B kinase subunit beta; I-kappa-B-kinase

Alternative Names beta; IKK-B; IKK-beta; IkBKB; I-kappa-B kinase 2; IKK2; Nuclear factor NF-kappa-B

inhibitor kinase beta; NFKBIKB

**Gene ID** 3551

SwissProt ID 014920.

## **Application**

**Dilution Ratio** WB: 1:500-1:1000 IP: 1:20

Molecular Weight Calculated MW: 87 kDa; Observed MW: 87 kDa

# Product Name: IKK beta Rabbit Monoclonal Antibody Catalog #: AMRe02148

**C** EnkiLife


Background

The NF-κB/Rel transcription factors are present in the cytosol in an inactive state, complexed with the inhibitory IκB proteins (1-3). Most agents that activate NF-κB do so through a common pathway based on phosphorylation-induced, proteasome-mediated degradation of IκB (3-7). The key regulatory step in this pathway involves activation of a high molecular weight IκB kinase (IKK) complex whose catalysis is generally carried out by three tightly associated IKK subunits.

### Research Area

**Signal Transduction** 

## **Image Data**



Western blot analysis of IKK beta in HUVEC lysates using IKK beta antibody.

#### Note

For research use only.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838