

## Summary

<b>Production Name</b>	Smad3 Rabbit Monoclonal Antibody
<b>Description</b>	Rabbit Monoclonal antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB,IP
<b>Reactivity</b>	Human,Mouse,Rat,Hamster

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Monoclonal
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA
<b>Purification</b>	Affinity Purification

## Immunogen

<b>Gene Name</b>	SMAD3
<b>Alternative Names</b>	SMAD3; MADH3; Mothers against decapentaplegic homolog 3; MAD homolog 3; Mad3; Mothers against DPP homolog 3; hMAD-3; JV15-2; SMAD family member 3; SMAD 3; Smad3; hSMAD3
<b>Gene ID</b>	4088
<b>SwissProt ID</b>	P84022.

## Application

<b>Dilution Ratio</b>	WB: 1:500-1:1000 IP: 1:20
<b>Molecular Weight</b>	Calculated MW: 48 kDa; Observed MW: 52 kDa

**Product Name: Smad3 Rabbit Monoclonal Antibody**  
**Catalog #: AMRe02613**




## Background

Smad3 transcription factor phosphorylated and activated by TGF-beta-type receptors. A receptor-regulated Smad (R-smad). Binds directly to consensus DNA-binding elements in the promoters of target genes. In mouse required for establishment of the mucosal immune response and proper development of skeleton.


## Research Area

Signal Transduction

## Image Data




Western blot analysis of Smad3 in 3T3, C6 lysates using Smad3 antibody.



Western blot analysis of Smad3 in Jurkat lysates using Smad3 antibody.

**Product Name: Smad3 Rabbit Monoclonal Antibody**  
**Catalog #: AMRe02613**



Western blot analysis of Smad3 in K562, CHO-K1 lysates using Smad3 antibody

### **Note**

For research use only.