

## Summary

<b>Production Name</b>	STING Rabbit Monoclonal Antibody
<b>Description</b>	Rabbit Monoclonal antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB,IHC-P
<b>Reactivity</b>	Human,Rat

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Monoclonal
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA
<b>Purification</b>	Affinity Purification

## Immunogen

<b>Gene Name</b>	STING1
<b>Alternative Names</b>	ERIS; MITA; MPYS; SAVI; NET23; STING; hMITA; hSTING; STING-beta;Transmembrane Protein 173
<b>Gene ID</b>	340061
<b>SwissProt ID</b>	Q86WV6.

## Application

<b>Dilution Ratio</b>	WB: 1:500-1:1000 IHC: 1:50-1:100
<b>Molecular Weight</b>	Calculated MW: 42 kDa; Observed MW: 42 kDa

**Product Name: STING Rabbit Monoclonal Antibody**  
**Catalog #: AMRe02694**




## Background

Facilitator of innate immune signaling that promotes the production of type I interferon (IFN-alpha and IFN-beta). Innate immune response is triggered in response to non-CpG double-stranded DNA from viruses and bacteria delivered to the cytoplasm.


## Research Area

Immunology

## Image Data



Western blot analysis of TMEM173 in K562, rat Brain lysates using Transmembrane Protein 173 antibody.



Immunohistochemistry analysis of paraffin-embedded Human tonsil using TMEM173 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

## Note

For research use only.