

Summary

Production Name	Phospho-beta Arrestin 1 (Ser412) Rabbit Monoclonal Antibody
Description	Rabbit Monoclonal antibody
Host	Rabbit
Application	WB,IHC-P,IP
Reactivity	Human

Performance

Conjugation	Unconjugated
Modification	Phosphorylated
lsotype	IgG
Clonality	Monoclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw
	cycles.
Buffer	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05%
	BSA
Purification	Affinity Purification

Immunogen

Gene Name	ARRB1
Alternative Names	ARB1; ARR1; ARRB1; ARRB1_HUMAN; Arrestin 2; Arrestin beta 1; Arrestin beta-1; Beta-
	arrestin-1.
Gene ID	408
SwissProt ID	P49407.

Application

Dilution Ratio	WB: 1:500-1:1000 IHC: 1:50-1:100 IP: 1:20
Molecular Weight	Calculated MW: 47 kDa; Observed MW: 50 kDa

Background

Members of arrestin/beta-arrestin protein family are thought to participate in agonist-mediated desensitization of Gprotein-coupled receptors and cause specific dampening of cellular responses to stimuli such as hormones, neurotransmitters, or sensory signals. Arrestin beta 1 is a cytosolic protein and acts as a cofactor in the beta-adrenergic receptor kinase (BARK) mediated desensitization of beta-adrenergic receptors. Besides the central nervous system, it is expressed at high levels in peripheral blood leukocytes, and thus the BARK/beta-arrestin system is believed to play a major role in regulating receptor-mediated immune functions. Alternatively spliced transcripts encoding different isoforms of arrestin beta 1 have been described. [provided by RefSeq, Jan 2011]

Research Area

Signal Transduction

Image Data



Western blot analysis of Phospho-beta Arrestin 1 (Ser412) in A549, MCF-7, U251, HL-60 lysates using Phospho-beta Arrestin



Immunohistochemistry analysis of paraffin-embedded Human lung cancer using beta Arrestin 1 (Phospho-Ser412) antibody.High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.



Note For research use only.