

Summary

| | |
|------------------------|----------------------------------|
| Production Name | IKZF3 Rabbit Monoclonal Antibody |
| Description | Rabbit Monoclonal antibody |
| Host | Rabbit |
| Application | WB,IHC-P |
| Reactivity | Human |

Performance

| | |
|---------------------|--|
| Conjugation | Unconjugated |
| Modification | Unmodified |
| Isotype | IgG |
| Clonality | Monoclonal |
| Form | Liquid |
| Storage | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. |
| Buffer | 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA |
| Purification | Affinity Purification |

Immunogen

| | |
|--------------------------|---|
| Gene Name | IKZF3 |
| Alternative Names | IKZF3; ZNFN1A3; Zinc finger protein Aiolos; Ikaros family zinc finger protein 3 |
| Gene ID | 22806 |
| SwissProt ID | Q9UKT9. |

Application

| | |
|-------------------------|--|
| Dilution Ratio | WB: 1:500-1:1000 IHC: 1:50-1:100 |
| Molecular Weight | Calculated MW: 58 kDa; Observed MW: 58 kDa |

Background

Product Name: IKZF3 Rabbit Monoclonal Antibody
Catalog #: AMRe03045




Transcription factor that plays an important role in the regulation of lymphocyte differentiation. Plays an essential role in regulation of B-cell differentiation, proliferation and maturation to an effector state. Involved in regulating BCL2 expression and controlling apoptosis in T-cells in an IL2-dependent manner.


Research Area

Epigenetics and Nuclear Signaling

Image Data



Western blot analysis of IKZF3 in HL-60, U251 lysates using IKZF3 antibody.



Immunohistochemistry analysis of paraffin-embedded Human tonsil using IKZF3 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

Note

For research use only.