

Summary

Production Name	Nampt Rabbit Monoclonal Antibody
Description	Rabbit Monoclonal antibody
Host	Rabbit
Application	WB,IHC-F,IHC-P,ICC/IF,IP
Reactivity	Human,Mouse,Rat

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Monoclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA
Purification	Affinity Purification

Immunogen

Gene Name	NAMPT
Alternative Names	Nicotinamide phosphoribosyltransferase; NAmPRTase; Nampt; Pre-B-cell colony-enhancing factor 1; Pre-B cell-enhancing factor; Visfatin
Gene ID	10135
SwissProt ID	P43490.

Application

Dilution Ratio	WB: 1:500-1:1000 IHC: 1:50-1:100 IF: 1:50-1:200 IP: 1:20
Molecular Weight	Calculated MW: 56 kDa; Observed MW: 56 kDa

Product Name: Nampt Rabbit Monoclonal Antibody
Catalog #: AMRe03816




Background

PBEF1: nicotinamide phosphoribosyltransferase. This gene encodes a protein that catalyzes the condensation of nicotinamide with 5-phosphoribosyl-1-pyrophosphate to yield nicotinamide mononucleotide, one step in the biosynthesis of nicotinamide adenine dinucleotide. The protein is an adipokine that is localized to the bloodstream and has various functions, including the promotion of vascular smooth muscle cell maturation and inhibition of neutrophil apoptosis. It also activates insulin receptor and has insulin-mimetic effects, lowering blood glucose and improving insulin sensitivity. The protein is highly expressed in visceral fat and serum levels of the protein correlate with obesity.


Research Area

Cardiovascular

Image Data




Immunocytochemistry analysis of Nampt (green) in U87-MG using Nampt antibody, and DAPI (blue).




Western blot analysis of Visfatin in 3T3, Raw264.7 lysates using Visfatin antibody.

Product Name: Nampt Rabbit Monoclonal Antibody
Catalog #: AMRe03816



Western blot analysis of Visfatin in HeLa, A549, HL-60, U2OS, C6 lysates using Visfatin antibody



Immunohistochemistry analysis of paraffin-embedded Human lung cancer using Visfatin antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

Note

For research use only.